

HEAVY METAL CONCENTRATIONS IN SOME ORGANS OF AFRICAN CATFISH  
(*Clarias gariepinus*) FROM EKO-ENDE DAM, IKIRUN, NIGERIA.

A. Adeyeye and P. B. Ayoola

Department of Science Laboratory Technology, Ladoke Akintola University of Technology, P. M.

B. 4000, Ogbomoso, Nigeria.

ABSTRACT

The contents of the heavy metals Fe, Zn, Cu, Mn, Pb, Cd and As were examined in the liver, heart, gills, kidney and muscles of the fish *Clarias gariepinus* from Eko-Ende dam in Ikirun, the capital of Ifelodun Local Government of Osun State, Nigeria. *C. gariepinus* is the fish of choice and the most demanded in the southwest of Nigeria. The highest metal concentrations were in the liver and the gills while the lowest was in the muscles. The general decreasing order of metal accumulation in the organs was Fe > Zn > Cu > Mn > Cd. Lead and arsenic were not detected in any organ. The values were of lower concentrations than found in many other dams and rivers in Nigeria and some other countries. The values were also lower than the FAO/WHO recommended maximum limits in fish samples, making the fish to be safe and not of any hazards for the consumers.

KEYWORDS: Heavy metals; organs; *Clarias gariepinus*; Eko-Ende dam; Nigeria.

Received for Publication: 03/11/12

Accepted for Publication: 24/01/13

Corresponding Author:

INTRODUCTION

The indispensable need for portable water has necessitated the need for construction of dams in various locations in Nigeria, and particularly in the southwest of the country, as reservoirs to supply portable water and thereby prevent water-borne diseases and improve the quality of life of the people. These dams also serve as sources of various species of fish for the populace. However, the contamination of fresh waters with a wide range of pollutants has become a matter of concern over the last few decades. Among the pollutants, the natural aquatic systems may extensively be contaminated with heavy metals released from domestic, industrial and other man-made activities, and through weathering of rocks and leaching of soils. Such metals may be precipitated, adsorbed on solid surface, remain soluble or suspended in water, or may be taken up by fauna and flora and eventually be accumulated in marine organisms including fish that are consumed by human beings (Adefemi *et al.*, 2008). Compared to other types of aquatic pollution, heavy metal pollution is less visible but its effects on the ecosystem and on man are intensive and extensive (Edem *et al.*, 2009). The fact that the toxic heavy metals are non-biodegradable and have the ability to accumulate in the environment makes them deleterious to the aquatic environment and consequently to man who depend on aquatic products including fish as sources of food. Fishes particularly are notorious for their ability to concentrate metals in their body tissues to dangerously poisonous levels (Kakulu & Osibanjo, 1988). The Minamata catastrophic disease of the 1950s and 1960s in Japan, caused by methyl mercury poisoning of people who consumed contaminated fish over a period of time is an example (Lars Jarup, 2003). Therefore, since they play an important role in human nutrition, fish need to be screened to ensure that unnecessarily high levels of the toxic heavy metals are not being transferred to man through them. African catfish (*Clarias gariepinus*) is of great commercial importance because it is the most widely consumed freshwater fish in Nigeria (Olaifa *et al.*, 2004). It is therefore a good choice to study its response to environmental contaminants, particularly the heavy metals. Eko-Ende dam is a major source of portable water to Ifelodun Local Government of Osun State, in the southwest of Nigeria, comprising of Ikirun and some other major towns and several other villages. It is also a veritable source of fish for the Local Government as well as the neighbouring Osogbo, the State capital. African catfish, *C. gariepinus*, is the most common and the most preferred fish from the dam. However, there has been a paucity of data concerning the heavy metal contents of



All rights reserved

This work by [Wilolud Journals](http://www.wiloludjournal.com) is licensed under a [Creative Commons Attribution 3.0 Unported License](http://creativecommons.org/licenses/by/3.0/)

the fishes from the dam. Therefore, this work provides information on the heavy metal concentrations in the organs of African catfish, *C. gariepinus*, in an attempt to predict the safety of its consumption in the environment. It could also be used to establish a baseline for future studies of heavy metal pollution in South Western Nigeria.

#### MATERIALS AND METHODS

Fresh adult samples of *C. gariepinus* were purchased from fishermen at Eko-Ende dam site and brought immediately into the laboratory and kept frozen prior to analysis. The fish were dissected into the various organs. The fish organs were oven-dried at 105°C to constant weight, blended into fine powder and each kept in a plastic container and stored in a refrigerator at about -4°C prior to analysis. Each separate organ was digested using freshly prepared concentrated nitric acid and hydrogen peroxide (1:1 v/v) according to FAO methods (FAO/SIDA, 1983). Reagent blanks were prepared accordingly to test the purity of the reagents, while standards of the metals in solution were used to calibrate the instrument. The metals were determined using the atomic absorption spectrophotometer (AAS), according to the AOAC (1995). The results were expressed as means of triplicate determinations. Data analysis was by one way analysis of variance (ANOVA) to assess the significance at  $p < 0.05$  using student t-test.

#### RESULTS AND DISCUSSION

The mean concentrations of the heavy metals (Fe, Zn, Cu, Mn, Pb, Cd and As) in the various organs of *C. gariepinus* in this study are shown in Table 1. The results confirmed the varying degrees of accumulation of the heavy metals by different organs. The highest heavy metal concentrations were found in the liver, followed by the gills and the least generally being in the muscles. The general distribution of the metals in the organs was in the order liver > gills > kidney > heart > muscle except for Mn with the order liver > gills > kidney > muscle > heart. The liver is a target organ for the accumulation of heavy metals, and the higher levels of these metals in it relative to other organs may be attributed to the high coordination of metallothionein protein with the metals (Hogstrand and Haux, 1991). In addition, the liver is the principal organ responsible for the detoxification, transportation and storage of toxic substances and an active site of pathological effects induced by contamination. The gills are the respiratory sites and are directly in contact with water and pollutants that may be present in

Table 1: Heavy metal concentrations (mg/kg dry weight) in organs of *Clarias gariepinus* samples from Eko-Ende dam.

| Metal | Liver      | Heart     | Gills      | Muscle    | Kidney     |
|-------|------------|-----------|------------|-----------|------------|
| Zn    | 1.92±0.18  | 1.13±0.06 | 1.45±0.16  | 0.80±0.05 | 1.39±0.18  |
| Cu    | 0.88±0.25  | 0.40±0.15 | 0.68±0.08  | 0.15±0.05 | 0.48±0.12  |
| Mn    | 0.55±0.18  | 0.18±0.04 | 0.45±0.06  | 0.24±0.04 | 0.28±0.03  |
| Fe    | 20.25±1.35 | 8.65±0.76 | 11.64±1.25 | 6.99±0.85 | 11.62±0.64 |
| Cd    | 0.03±0.01  | 0.03±0.01 | 0.02±0.01  | ND        | 0.01±0.01  |
| Pb    | ND         | ND        | ND         | ND        | ND         |
| As    | ND         | ND        | ND         | ND        | ND         |

Values are means (±SD) of triplicate determinations. ND = Not detected.

water. Thus, the concentrations of heavy metals in gills are a reflection of the concentration in the surrounding water. The high metal concentrations in the gills could be due to the metal complexing with the mucus that is impossible to remove completely from the lamellae before analysis (Heath, 1987).

Our results gave higher levels of heavy metals in the liver and gills relative to other organs. This is in agreement with the findings of others elsewhere (Zyadah, 1999; Farombi *et al.*, 2007; Dimari *et al.*, 2008; Al-weher, 2008; Yilmaz, 2009; Uzairu *et al.*, 2009; Akan *et al.*, 2009) (see also Table 2). Since the liver is a target organ of accumulation for many metals, it is often considered a good monitor of water pollution with heavy metals since their concentrations have proportional relevance to those present in the environment (Yilmaz, 2009). The muscles had the lowest metal concentrations compared to other organs. This agrees with other peoples' findings



All rights reserved

This work by Wilolud Journals is licensed under a [Creative Commons Attribution 3.0 Unported License](https://creativecommons.org/licenses/by/3.0/)

in some other places (Canli & Kalay, 1998; Ekpo *et al.*, 2008; Edem *et al.*, 2009; Yilmaz, 2009), showing that muscle is not an active tissue in accumulating heavy metals.

Iron was the most abundant of all the metals considered in this study, and Zn the second most abundant. The general decreasing order of metal accumulation in the organs (liver, gills, kidney and heart) was  $Fe > Zn > Cu > Mn > Cd$  except in the muscles which order was  $Fe > Zn > Mn > Cu > Cd$ . Lead and arsenic were not detected in any organ. It is generally accepted that heavy metal uptake occurs mainly from water, food and sediment. However, the efficiency may differ in relation to ecological needs, metabolism and the contamination gradients of water, food and sediment, as well as other factors such as salinity, temperature and interacting agents (Pagenkopf, 1983).

The higher levels of Fe and Zn in the fish organs relative to other metals could be due to the fact that these metals are naturally abundant in Nigerian soils which is the main source of metals in the surrounding water of the fish samples (Adefemi *et al.*, 2008); Kakulu & Osibanjo, 1988). In general, accumulation of the essential elements Zn, Cu, Fe and Mn were higher than the non-essential Pb, Cd and As in the fish organs. The essential elements play vital biochemical and physiological functions in fish. Zinc, for example, is regulated to maintain a certain homeostatic status in fish (Chen & Chen, 1999), while both Fe and Cu are components of the enzyme cytochrome oxidase which is involved in energy metabolism (NAS, 1976). But when in excess of the body needs of fish or man, these metals may constitute a major pollution source and pose a serious health risk (Onyia *et al.*, 2007). Toxicity of iron, for example, may lead to haemochromatosis and, in severe cases, to thalassaemia (Hovinga *et al.*, 1993). Excessive intake of zinc may lead to diarrhea and vomiting, while toxicity to manganese leads to a syndrome called manganism which involves both psychiatric symptoms and features of Parkinson disease (Dobson *et al.*, 1994).

The non-essential heavy metals, such as lead and cadmium, have no biological function, but are rather detrimental to fish and human existence even at very low concentrations. Acute toxicity of cadmium, for example, may lead to high blood pressure, red blood cells destruction and kidney and testicular tissue destruction, while lead toxicity can reduce intelligence, delay motor development, impair memory and cause hearing problems (Sharma *et al.*, 2008).

From the results of this study, it is seen that fish from Eko-Ende dam is less polluted than those from many other locations in Nigeria, including those from Ogun River in the southwest, Lake Chad, Kubani River in Zaria, Alau dam in Maiduguri and the Niger Delta, (Table 2). The values were also lower than those from some other countries, such as from Jordan, Taiwan and Turkey (Table 2). This could be attributed to the location of the dam in a purely agricultural environment devoid of any major industrial activities.

In conclusion, the heavy metals in this study showed an uneven distribution in the organs of the fish, with the metals being more concentrated in the liver and the gills. They had their concentrations far below the FAO/WHO and IAEA maximum permissible levels (Table 2). With these findings, one may admit that the levels of the metals in the fish samples would be safe and not constitute any health hazards for the consumers. It is however important that continuous monitoring exercise be put in place to guide against excessive bioaccumulation of these metals and safeguard the safety, protection and well-being of the consumers.



All rights reserved

This work by [Wilolud Journals](#) is licensed under a [Creative Commons Attribution 3.0 Unported License](#)

Table 2. Heavy metal concentrations in fish (mg/kg) in studies from some other locations in Nigeria and other countries.

| Location/ (Fish)                | Organ | Zn    | Cu    | Mn   | Fe                   | Cd    | Pb     | Reference                       |                          |      |       |      |       |        |       |       |  |
|---------------------------------|-------|-------|-------|------|----------------------|-------|--------|---------------------------------|--------------------------|------|-------|------|-------|--------|-------|-------|--|
| Ogun River, Nigeria             | Liver | 19.75 | 4.70  | --   | --                   | 2.10  | 3.40   | Farombi <i>et al.</i> , 2007    | (Clarias gariepinus)     |      |       |      | Gill  | 20.35  | 4.55  | --    |  |
| -- 2.05 2.40                    |       |       |       |      |                      |       | kidney | 19.05 5.00 --                   | --                       | --   | 0.69  | 3.35 |       |        |       |       |  |
| Alau Dam, Maiduguri, Nigeria    | Heart | 8.96  | 2.19  | --   | --                   | 0.25  | 1.69   |                                 |                          |      |       |      |       |        |       |       |  |
| 0.62 0.33 0.38                  | Liver | 0.52  | 0.44  | 0.35 | 0.40                 | 0.49  | 0.40   | Dimari <i>et al.</i> , 2008     | (Tilapia gallier)        |      |       |      |       | Gill   | 0.36  | 0.63  |  |
|                                 | 0.53  |       |       |      |                      |       |        | Intestine                       | 0.15                     | 0.36 | 0.14  | 0.34 | 0.09  | 0.12   |       |       |  |
| Okumeshi River, Delta State     | Liver | --    | --    | 0.49 | --                   | 0.31  | 0.01   | Ekeanyanwu <i>et al.</i> , 2010 | (Tilapia)                |      |       |      |       | Gill   | --    | --    |  |
| 0.17 -- 0.21                    | <0.01 |       |       |      |                      |       |        | Muscle                          | --                       | --   | 1.97  | --   | 0.62  | <0.01  |       |       |  |
| Kubani River, Zaria, Nigeria    | Liver | 49.56 | 19.31 | --   | --                   | 0.21  | 0.28   | Uzairu <i>et al.</i> , 2009     | a. (Clarias gariepinus)  |      |       |      |       | Gill   | 20.05 | 1.87  |  |
| -- -- 0.31                      | 0.12  |       |       |      |                      |       |        | Muscle                          | 4.92                     | 0.24 | --    | --   | 0.02  | 0.04   |       |       |  |
| b. (Oreochromis niloticus)      | Liver | 65.72 | 40.11 | --   | --                   | 0.40  | 0.76   |                                 |                          |      |       |      |       | Gill   | 26.14 | 5.32  |  |
| -- -- 0.08                      | 0.24  |       |       |      |                      |       |        | Muscle                          | 7.61                     | 1.15 | --    | --   | 0.02  | 0.02   |       |       |  |
| Lake Chad, Nigeria              | Liver | 0.54  | 0.52  | 0.32 | 0.34                 | 0.22  | 0.32   | Akan <i>et al.</i> , 2009       | a. (Tilapia zilli)       |      |       |      |       | Gill   | 0.33  | 0.45  |  |
| 0.29 0.28 0.01                  | 0.03  |       |       |      |                      |       |        | Kidney                          | 0.12                     | 0.31 | 0.13  | 0.15 | 0.10  | 0.01   |       |       |  |
| b. (Clarias anguillaris)        | Liver | 0.31  | 0.32  | 0.29 | 0.26                 | 0.03  | 0.15   |                                 |                          |      |       |      |       | Gill   | 0.28  | 0.32  |  |
| 0.22 0.24 0.02                  | 0.10  |       |       |      |                      |       |        | Kidney                          | 0.13                     | 0.21 | 0.11  | 0.12 | 0.01  | 0.02   |       |       |  |
| Henshaw Beach, Calabar, Nigeria | Liver | 0.257 | --    | --   | --                   | 0.049 | 0.173  | Edem <i>et al.</i> , 2009       | (Oreochromis niloticus)  |      |       |      |       | Gills  | 0.198 | --    |  |
| -- 0.038 0.133                  |       |       |       |      | (group A: 29cm size) |       |        | Muscles                         | 0.079                    | --   | --    | --   | 0.015 | 0.053  |       |       |  |
| Northern Jordan Valley, Jordan  | Gill  | 40.12 | 9.12  | --   | --                   | 0.77  | --     | Al-Weher, 2008                  | (Clarias lazera)         |      |       |      |       | Muscle | 30.13 | 3.04  |  |
| -- -- 0.24                      | --    |       |       |      |                      |       |        |                                 |                          |      |       |      |       |        |       |       |  |
| Köycegiz Lake–Mugla, Turkey.    | Liver | 136.9 | 114.3 | 9.64 | --                   | 0.44  | 1.02   | Yilmaz, 2009                    | (Oreochromis niloticus)  |      |       |      |       | Gill   | 104.8 | 7.34  |  |
| -- 0.11 0.96                    |       |       |       |      |                      |       |        | Muscle                          | 84.76                    | 3.91 | 12.65 | --   | 0.12  | 1.12   |       | 63.44 |  |
| Coastal Waters, Eastern Taiwan  | Liver | 33.83 | 5.58  | --   | --                   | 0.52  | 0.21   | Wen-Bin Huang, 2003             | (mean for n = 81 fishes) |      |       |      |       | Gill   | 18.12 | 1.45  |  |
| -- -- 0.27                      | 0.32  |       |       |      |                      |       |        | Muscle                          | 3.79                     | 0.34 | --    | --   | 0.05  | 0.05   |       |       |  |
| FAO/WHO Limits                  |       | 40    | 30    |      |                      | 0.5   | 0.5    | FAO/WHO, 1989.                  |                          |      |       |      |       |        |       |       |  |
| IAEA – 407                      |       |       | 3.28  | 11.0 | 146                  | 0.18  | 0.12   | Wyse                            | et                       |      |       |      | al.,  |        |       | 2005  |  |



All rights reserved

This work by [Wilolud Journals](http://www.wiloludjournals.com) is licensed under a [Creative Commons Attribution 3.0 Unported License](http://creativecommons.org/licenses/by/3.0/)

## REFERENCES

- Abu Hilal, A. H. and Ismail, N. S. (2008). Heavy metals in eleven common species of fish from the Gulf of Aqaba, Red Sea. *Jordan J. Biol. Sci.* 1(1): 13-18.
- Adefemi, S. O., Asaolu, S. S. and Olaofe, O. (2008). Determination of heavy metals in *Tilapia mossambicus* fish, associated water and sediment from Ureje dam in South-Western Nigeria. *Res. J. Environ. Sci.* 2: 151-155.
- Akan, J. C., Abdulrahman, F. I., Sodipo, O. A. and Akandu, P. I. (2009). Bioaccumulation of some heavy metals of six fresh water fishes caught from Lake Chad in Doron Buhari, Maiduguri, Borno State, Nigeria. *J. Appl. Sci. Environ. Sanitation* 4(2): 103-114.
- Al-Weher, S. M. (2008). Levels of heavy metals Cd, Cu and Zn in three fish species collected from the Northern Jordan Valley, Jordan. *Jordan J. Biol. Sci.* 1(1): 41-46.
- AOAC (1995). *Official Methods of Analysis*. 16<sup>th</sup> Edn. Association of Official Analytical Chemists, Washington D C.
- Canli, M. and Kalay, Ö. AY, M. (1998). Levels of heavy metals (Cd, Pb, Cu, Cr and Ni) in tissues of *Cyprinus carpio*, *Barbus capito* and *Chondrostoma regium* from the Seyhan River, Turkey. *Tr. J. Zoology* 22: 149-157.
- Chen, M. H. and Chen, C. Y. (1999). Bioaccumulation of sediment-bound heavy metals in grey mullet, *Liza macrolepis*. *Mar. Pollut. Bull.* 39: 239-244.
- Dimari G. A., Abdulkarim F. I., Akan J. C. and Garba S. T. (2008). Metal concentrations in tissues of *Tilapia galier*, *Clarias lazera* and *Osteoglosidae* caught Alau dam, Maiduguri, Bornu State, Nigeria. *American J. Environ. Sci.* 4(4): 373-379.
- Dobson A. W., Erickson K. M. and Aschner M. (2004). Manganese neurotoxicity. *Ann. NY Acad. Sci.*, 1012: 115-128.
- Edem, C. A., Osabor, V., Iniama, G., Etiuma, R. and Eke, J. (2009). Distribution of heavy metals in bones, gills, liver and muscles of (Tilapia) *Oreochromis niloticus* from Henshaw Town Beach market, in Calabar, Nigeria. *Pak. J. Nutrition*, 8(8): 1209-1211.
- Ekeanyanwu, C. R., Ogbuinyi, C. A. and Etienajirhevwe, O. F. (2010). Trace metals distribution in fish tissues, bottom sediments and water from Okumeshi River in Delta State, Nigeria. *Ethiopian J. Environ. Studies and Management*, 3(3): 12-17.
- Ekpo, K. E., Asia, I. O., Amayo, K. O. and Jegede, D. A. (2008). Determination of lead, cadmium and mercury in surrounding water and organs of some species of fish from Ikpoba River in Benin City, Nigeria. *Int. J. Phy. Sci.* 3(11): 289-292.
- FAO/SIDA (1983). Manual of methods in aquatic environment research. Part 9. Analyses of metals and organochlorines in fish. *FAO Fish Tech Paper* 212. 33pp.
- FAO/WHO (1989). Evaluation of certain food additives and the contaminants mercury, lead and cadmium. *WHO Tech. Report*. Series No. 505.
- Farombi, E. O., Adelowo, O. A. and Ajimoko, Y. R. (2007). Biomarkers of oxidative stress and heavy metal levels as indicators of environmental pollution in African catfish (*Clarias gariepinus*) from Nigeria Ogun River. *Int. J. Environ. Res. Public Health*, 4(2): 158-165.
- Heath, A. G. (1987). *Water pollution and fish physiology*. CRC Press, Florida.



All rights reserved

This work by [Wilolud Journals](#) is licensed under a [Creative Commons Attribution 3.0 Unported License](#)

- Hogstrand, C. and Haux, C. (1991). Binding and detoxification of heavy metals in lower vertebrates with reference to metallothionein. *Comp. Biochem. Physiol.* 100 (1&2): 137-141.
- Hovinga M. E., Sowers M. and Humphrey H. E. (1993). Environmental exposure and lifestyle predictors of lead, cadmium and PCB and DDT levels in Great Lakes fish eaters. *Arch. Environ. Health*, 48: 98-104.
- Kakulu, S. E. and Osibanjo, O. (1988). Trace heavy metal pollution status in sediments of Niger Delta area. *J. Chem. Soc. Nigeria*, 13: 9-11.
- Lars Jarup. (2003). Hazards of heavy metal contamination. *British Med. Bull.* 68: 167- 182.
- NAS (1976). Food and Nutrition Board. Zinc in human nutrition. In: *Introduction to Nutrition*, Fleck, H. (Ed). 3<sup>rd</sup> Edn. Macmillan, New York.
- Olaifa, F. E., Olaifa, A. K. and Onwude, T. E. (2004). Lethal and sublethal effects of copper to the African catfish (*Clarias gariepinus*). *Afri. J. Biomed. Res.* 7: 65-70.
- Onyia L. U., Millam C. and Ese D. S. (2007). Investigation of heavy metals in four commercial fishes along Upper River Benue, Yola. *Inter. J. Physical Sci.* 2(2):
- Pagenkopf, G. K. (1983). Gill surface interaction model for trace metal toxicity of fish. Role of complexation, pH, water hardness. *Environ. Sci. Technol.* 17(6): 342-347.
- Sharma J. L., Garg N. K. and Buldini P. L. (2008). *Condensed Chemical Dictionary*. CBS Publishers, New Delhi, India.
- Uzairu, A., Harrison, G. F. S., Balarabe, M. L. and Nnaji, J. C. (2009). Concentration levels of trace metals in fish and sediment from Kubani River, Northern Nigeria. *Bull. Chem. Soc. Ethiopia*, 23(1): 9-17.
- Wen- Bin Huag (2003). Heavy metal concentrations in the common Benthic fishes caught from the coastal waters of Eastern Taiwan. *J. Food Drug Analysis* 11(4): 324-330.
- Wyse, E. J., Azemard, S. and Mora, S. J. (2005). *Report on the world-wide inter-comparison exercise for the determination of trace elements and methyl mercury in fish homogenate*. IAEA-407, IAEA/AL/144 (IAEA/MEL/72), IAEA, Monaco.
- Yilmaz, F. (2009). The comparison of heavy metal concentrations (Cd, Cu, Mn, Pb and Zn) in tissues of three economically important fish (*Anguilla anguilla*, *Mugil cephalus* and *Oreochromis niloticus*) inhabiting Köycegiz Lake-Mugla (Turkey). *Turkish J. Sci. Technol.* 4(1): 7-15.
- Zyadah, M. A. (1999). Accumulation of some heavy metals in *Tilapia zilli* organs from Lake Manzalah, Egypt. *Tr. J. Zoology* 23: 365-372.
- 



All rights reserved

This work by [Wilolud Journals](#) is licensed under a [Creative Commons Attribution 3.0 Unported License](#)